



Press Release 16-089

Earlier snowmelt decreases streamflow, reduces forests' ability to regulate atmospheric carbon dioxide

By mid-century, shift in snowmelt timing could lead to 45 percent reduction of forest CO2 uptake



A Colorado Rocky Mountain forest during the winter-spring transition period, which is changing.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)

August 3, 2016

Earlier annual snowmelt periods may hinder the ability of forests to regulate atmospheric carbon dioxide (CO₂), according to the results of a new study.

The findings, published in *Geophysical Research Letters* <<https://www.nsf.gov/cgi-bin/good-bye?http://onlinelibrary.wiley.com/doi/10.1002/2016GL069769/abstract>>, a journal of the American Geophysical Union, predict that this shift in snowmelt timing each spring could result in a 45 percent reduction of snowmelt period forest carbon by mid-century.

A second study, also published <<https://www.nsf.gov/cgi-bin/good-bye?http://onlinelibrary.wiley.com/doi/10.1002/2016GL069690/abstract>> in *Geophysical Research Letters*, found that earlier, slower snowmelt reduces the amount of streamflow, which has consequences for agriculture, municipal water supplies and recreational opportunities in Colorado and other states in the western U.S.

"The recent western drought has been accompanied by a snowpack restricted to higher elevations, with a significant effect on the ski industry," said Tom Torgersen, program director in the National Science Foundation (NSF) Division of Earth Sciences, which funded the research. NSF's Long-Term Ecological Research (LTER) program also supported the studies through the Niwot Ridge, Colorado, LTER site.



Mid-summer snowfields in the mountains of Washington State; most snow has melted.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)



The Collegiate Peaks viewed from the Arkansas River Valley in Colorado.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)



Scientists gauge streamflow in the Sagehen Experimental Watershed in California.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)



Colorado's San Juan Mountains covered in late spring snow. Snowmelt timing is shifting here.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)

"Climate variability also leads to conditions favoring earlier and slower snowmelt, with a decreased and prolonged peak streamflow," Torgersen said. "This water flow affects mountain fishing and results in less forest growth. The effects of drought and climate variability on snowmelt reach far beyond farm productivity and urban water restrictions."



The Yellowstone River in Yellowstone National Park after a late spring snowstorm.

[Credit and Larger Version \(/news/news_images.jsp?cntn_id=189304&org=NSF\)](#)

Implications for western U.S.

Forests in seasonally snow-covered areas serve as key CO₂ sinks, thanks to the natural processes by which trees take in carbon. This carbon uptake is restrained during winter, but increases to peak capacity in spring when snowmelt provides abundant water to trees.

University of Colorado Boulder (CU-Boulder) scientists working at Niwot Ridge in Colorado's Rocky Mountains studied 15 years of snowmelt and atmospheric CO₂ data to determine the effects of changes in snowmelt periods.

They found that earlier snowmelt triggered by climate change reduces forests' ability to take CO₂ out of the atmosphere.

"The implications of this research are profound as mountains in the western U.S. are an important part of the regional cycling of carbon and water," said Noah Molotch, the director of CU-Boulder's Center for Water, Earth Science & Technology, and a co-author of both studies.

Added Taylor Winchell of CU-Boulder's Institute for Arctic and Alpine Research (INSTAAR) and lead author of one of the studies: "Early melting reduces trees' ability to uptake carbon during the snowmelt period, a key time for seasonal carbon uptake."

Downstream water resources

Snowmelt also provides water resources to downstream communities. Previous research shows that the timing and rate at which snow melts can affect the amount and quality of water available for vegetation, farming and fishing.

The researchers used a unique modeling system to study the effects of earlier snowmelt across various regions of the western U.S., including the Cascade Range, the Sierra Nevada, the Wasatch Range and the Rocky Mountains. These areas see significant seasonal snow accumulations that generate water resources for downstream communities.

The results show that earlier, slower snowmelt, triggered by warmer temperatures, reduces streamflow. These slower "trickle" melts reduce percolation in hillslope soils and allow more water to evaporate, resulting in less streamflow overall.

"Of all the regions we studied, streamflow from Colorado's Rocky Mountains is most sensitive to changes in snowmelt," said Theodore Barnhart of INSTAAR, lead author of the second study. "This analysis suggests that all the regions studied will experience a decrease in streamflow with a decrease in snowmelt rate, with some regions having more streamflow sensitivity than others."

CU-Boulder's Molotch added that the findings have broad implications for the scientific community.

"Given that 60 million people in the western U.S. depend on snowmelt for their water supply, the future decline in snowmelt-derived streamflow may place additional stress on over-allocated water supplies," he said. "These two studies are reshaping the way scientists -- and land and water managers -- think about climate change in mountain regions."

-NSF-

Media Contacts

Cheryl Dybas, NSF, (703) 292-7734, cdybas@nsf.gov (<mailto:cdybas@nsf.gov>)

Trent Knoss, University of Colorado Boulder, (303) 735-0528, trent.knoss@colorado.edu (<mailto:trent.knoss@colorado.edu>)

Lauren Lipuma, American Geophysical Union, (202) 777-7396, llipuma@agu.org (<mailto:llipuma@agu.org>)

Related Websites

NSF Grant: Snowpack energy and mass balance: implications for biogeochemical feedbacks in alpine basins: https://www.nsf.gov/awardsearch/showAward?AWD_ID=1032308&HistoricalAwards=false <https://www.nsf.gov/awardsearch/showAward?AWD_ID=1032308&HistoricalAwards=false>

Study Ties Forest "Greenness" in Western U.S. to Snowpack Extent: https://www.nsf.gov/news/news_summ.jsp?cntn_id=125359&org=NSF <https://www.nsf.gov/news/news_summ.jsp?cntn_id=125359&org=NSF>

The National Science Foundation (NSF) is an independent federal agency that supports fundamental research and education across all fields of science and engineering. In fiscal year (FY) 2016, its budget is \$7.5 billion. NSF funds reach all 50 states through grants to nearly 2,000 colleges, universities and other institutions. Each year, NSF receives more than 48,000 competitive proposals for funding and makes about 12,000 new funding awards. NSF also awards about \$626 million in professional and service contracts yearly.

[Get News Updates by Email](http://service.govdelivery.com/service/subscribe.html?code=USNSF_51) <http://service.govdelivery.com/service/subscribe.html?code=USNSF_51>

Useful NSF Web Sites:

NSF Home Page: <https://www.nsf.gov> <<https://www.nsf.gov>>

NSF News: [https://www.nsf.gov/news/ \(/news/\)](https://www.nsf.gov/news/)

For the News Media: [https://www.nsf.gov/news/newsroom.jsp \(/news/newsroom.jsp\)](https://www.nsf.gov/news/newsroom.jsp)

Science and Engineering Statistics: [https://www.nsf.gov/statistics/ \(/statistics/\)](https://www.nsf.gov/statistics/)

Awards Searches: [https://www.nsf.gov/awardsearch/ \(/awardsearch/\)](https://www.nsf.gov/awardsearch/)

National Science Foundation, 4201 Wilson Boulevard, Arlington, Virginia 22230, USA Tel: (703) 292-5111, FIRS: (800) 877-8339 | TDD: (800) 281-8749